

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 25/30, 57/20, 43/40, 47/36, 43/50	A1	(11) International Publication Number: WO 00/42847 (43) International Publication Date: 27 July 2000 (27.07.00)
(21) International Application Number: PCT/US00/00945 (22) International Filing Date: 14 January 2000 (14.01.00) (30) Priority Data: 09/233,796 19 January 1999 (19.01.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/233,796 (CON) Filed on 19 January 1999 (19.01.99) (71) Applicant (for all designated States except US): AMWAY CORPORATION [-/US]; 7575 Fulton Street, Ada, MI 49355 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): BRUMBAUGH, Ernest, H. [US/US]; 11029 Crooked Lake Road, Rockford, MI 49341 (US). (74) Agent: CONFORTI, Vita, G.; Brinks Hofer Gilson & Lione, P.O. Box 10087, Chicago, IL 60610 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: PLANT PROTECTING ADJUVANT CONTAINING TOPPED OR PEAKED ALCOHOL ALKOXYLATES AND CONVENTIONAL ALCOHOL ALKOXYLATES (57) Abstract A plant protecting composition comprising a biocide component and an adjuvant component wherein the adjuvant component includes at least one surfactant selected from the group consisting of topped alcohol alkoxylates and peaked alcohol alkoxylates and combinations thereof in combination with a conventional alcohol alkoxylate.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PLANT PROTECTING ADJUVANT CONTAINING TOPPED OR PEAKED ALCOHOL ALKOXYLATES AND CONVENTIONAL ALCOHOL ALKOXYLATES

BACKGROUND OF THE INVENTION

5 This invention relates to plant protecting compositions containing adjuvant compositions, and in particular, to adjuvant compositions that contain a topped or peaked alcohol ethoxylate or combinations thereof in combination with a conventional alcohol ethoxylate.

10 Adjuvants are commonly used in plant protecting compositions to facilitate biocidal action or to facilitate or modify characteristics of the biocidal formulations or spray solutions. In order to enhance or modify the chemical and/or physical characteristics of certain pesticides, adjuvants are added to form a mixture for spraying. Since spray application can be critical to the performance of the agricultural chemical, adjuvants are added to reduce
15 application problems such as chemical stability, incompatibility, solubility, suspension, foaming, drift, evaporation, volatilization, phytotoxicity, surface tension, droplet size and coverage. They can, depending on their type, enhance wetting, translocation, spreading, sticking, emulsifying, dispersing and biological activity. Adjuvants include wetting agents, crop oil
20 concentrates, spreaders, stickers, buffering agents, foaming and anti-foaming agents, dispersing agents and drift control agents. Over 200 EPA-registered pesticides have specific recommendations on their labels for adjuvant use. For example, they are recommended to enhance biological activity of the pesticide and to reduce, minimize or eliminate spray application problems as
25 previously noted. There are several different types of adjuvants recommended. To achieve consistent and effective results, the user must first select the desired type of adjuvant and then the appropriate product within that specific type for use with a particular biocide and then use that product at recommended rates. Adjuvants can be added to the biocide formulation
30 (commonly called "ready-to-dilute") or at the applicator stage when the

pesticide formulation is diluted with water or a suitable carrier into a spray tank (commonly called "tank mix").

Nonionic surfactants have been used as activator adjuvants to facilitate biocide penetration wetting, spreading, dispersing, solubilizing, emulsifying and other surface modifying properties to bring about enhanced biocidal action. These surfactants have also been used as spray-modifier adjuvants to facilitate sticking or spreading of wettable biocidal powders or to modify the drift properties of the biocide. Examples of common nonionic surfactants used as spray adjuvants can be found in Adjuvants for Herbicides (Weed Science Society of America 1982), which is incorporated herein by reference.

A common nonionic surfactant used as an adjuvant is an alkyl aryl oxylate such as nonyl ethoxylate phenol, also known as nonoxynol-10. Although nonoxynol-10 is effective, suitable alternatives are continually being sought.

An acceptable alternative is a conventional aliphatic alcohol alkoxyate, commonly called an alcohol alkoxyate, such as alcohol ethoxylates. Alcohol ethoxylates or ethylene oxide adducts are nonionic surfactants having functional properties such as wetting, foaming, emulsifying, and dispersing abilities as well as solubilization and detergent abilities. While the use of alcohol ethoxylates alone as adjuvants has been satisfactory, there is an ongoing effort to make adjuvants with improved herbicidal performance.

Surprisingly, it has been found that the addition of an adjuvant composition containing topped or peaked alcohol alkoxyates, or combinations thereof in combination with conventional alcohol alkoxyates to certain post emergent biocidal mixtures increases the efficacy of these biocides beyond that of recognized standards.

SUMMARY OF THE INVENTION

The present invention relates to a plant protecting composition comprising an effective amount of agricultural biocide and from about 1% to about 40% of an adjuvant composition comprising a combination of nonionic

5 surfactants selected from the group consisting of topped alcohol alkoxylates and peaked alcohol alkoxylates and combinations thereof with conventional alcohol alkoxylates. Preferably, the topped and peaked alcohol alkoxylates include alcohol ethoxylates, alcohol propoxylates, propoxylated and ethoxylated alcohols and combinations thereof. More preferably, the adjuvant composition includes primary aliphatic alcohol alkoxylates. The plant protecting composition may also contain other surfactants and additives so long as they do not detract from the improved results obtained from the adjuvant of the present invention.

10 The present invention further includes a method for increasing the efficacy of an agricultural biocide formulation comprising the step of providing an adjuvant comprising a nonionic surfactant selected from the group consisting of topped alcohol alkoxylates and peaked alcohol alkoxylates and combinations thereof with conventional alcohol alkoxylates. In one embodiment, the adjuvant composition and a biocide component are added to a tank mix. In another embodiment, the adjuvant is combined with the biocide component in a ready-to-dilute formulation.

5 Unexpectedly, it has been found that the adjuvant of the present invention improves the efficacy of certain post emergent agricultural biocide formulations beyond that of recognized standards.

0 It is noted that, unless otherwise stated, all percentages given in this specification and the appended claims refer to percentages by weight.

5 These and other advantages and features of the present invention will be better understood upon review of the following detailed description of the preferred embodiments.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention relates to a plant protecting composition containing an adjuvant composition comprising a nonionic surfactant component selected from the group consisting of topped alcohol ethoxylates

and peaked alcohol ethoxylates and combinations thereof with conventional alcohol alkoxyates.

Alcohol alkoxyates useful in the present invention are those containing an aliphatic hydrocarbon chain and an polyoxyalkylene chain.

5 The hydrocarbon chain may be derived from an aliphatic alcohol. Examples of alcohol alkoxyates useful as nonionic surfactants usually include those containing a fatty alcohol chain having a carbon length of from about C₈ to about C₂₀. The oxyalkylene chain of the compound can have a length of from C₂ to about C₁₀. Alcohol alkoxyates useful as nonionic surfactants will
10 preferably contain a oxyalkylene chain consisting mainly of ethylene oxide units (commonly called alcohol ethoxylates), propylene oxide units (commonly called alcohol propoxylates) or a combination thereof (commonly called propoxylated and ethoxylated alcohols).

15 Alcohol alkoxyates are generally prepared by alkoxyating the aliphatic alcohol with the oxyalkylene in the presence of a catalyst such as potassium oxide or sodium oxide. Examples of alcohol ethoxylates and alcohol propoxylates useful as nonionic surfactants include C₈-C₁₈ alcohols with 1-15 moles of ethylene oxide (EO) or propylene oxide (PO) units per mole of alcohol. The distribution of ethoxylation or propoxylation, as the case may be,
20 is quite broad and a sizable amount of free alcohol is left in the product. Common conventional alcohol ethoxylates are listed under the chemical classification of "ethoxylated alcohols" in McCutcheon's Emulsifiers & Detergents, Annual 1992. Common conventional alcohol propoxylates as well as propoxylated and ethoxylated alcohols are listed under the chemical
25 classification "propoxylated & ethoxylated fatty acids, alcohol or alkyl phenols" in McCutcheon's. The relevant portions of McCutcheon's are incorporated herein by reference.

30 Certain alcohol alkoxyates useful in the present invention are selected from the group consisting of topped alcohol alkoxyates, peaked alcohol alkoxyates and combinations thereof.

Topped alcohol alkoxylates refer to a distribution containing C₆ or higher carbon chain monofunctional and polyfunctional alcohols. Topped alcohol alkoxylates include both linear and branched alcohol alkoxylates. Topped alcohol alkoxylates have free alcohol and low mole alkoxylate entities removed physically by conventional separation processes such as extraction and distillation. Particularly preferred topped alcohol alkoxylates contain primary alcohol alkoxylates having C₆ or greater carbon chains.

Peaked alkoxylates refers to a narrow, high mole adduct distribution from the ethoxylation of linear or branched alcohols. Peaked alkoxylates have a more narrow and highly peaked alkoxylation distribution that results in a low amount of residual free alcohol, a lower amount of lower oxyalkylene adducts and a lower amount of higher oxyalkylene adducts in the product. Peaked alcohol alkoxylates are obtained through the use of different catalysts and/or manufacturing conditions. Examples of the preparation of peaked alcohol ethoxylates include U.S. Pat. No. 4,210,764 to Yang et al. and U.S. Pat. No. 5,118,650 to King, the disclosures of which are incorporated herein by reference.

The preferred peaked alcohol alkoxylates of the present invention include alcohol alkoxylates having a residual free alcohol content of less than about three percent. Particularly preferred peaked alcohol alkoxylates are C₆-C₂₀ alcohol ethoxylates, C₆-C₂₀ alcohol propoxylates, C₆-C₂₀ propoxylated and ethoxylated alcohols and combinations thereof. The more preferred peaked alcohol alkoxylates are C₈-C₁₆ alcohol ethoxylates containing from about 2 to about 20 moles of ethylene oxide (EO) per molecule, C₈-C₁₆ alcohol propoxylates containing from about 2 to about 20 moles of propylene oxide (PO) per molecule, C₈-C₁₆ propoxylated and ethoxylated alcohols.

The more preferred topped and peaked alcohol alkoxylates are the biodegradable aliphatic alcohol alkoxylates, in particular primary aliphatic alcohol alkoxylates, and more preferably linear primary alcohol alkoxylates.

The adjuvant composition of the present invention may also contain other ingredients found in commercial adjuvants so long as they do not

detract from the beneficial effects of the present invention. Examples of optional ingredients include additional surfactants, viscosity modifiers, stabilizing agents, pH modifiers, corrosion inhibitors, and the like. Examples of such optional ingredients can be found in U.S. Pat. No. 5,084,087 to Hazen et al., which is incorporated herein by reference.

The adjuvant composition of the present invention may be combined with one or more biocide components in various forms. For example, the adjuvant composition can be added to tank mixes with the biocide component(s). Alternatively, the adjuvant composition of the subject invention can be incorporated into a ready-to-dilute formulation containing the biocide component(s). Common forms of ready-to-dilute formulation include, for example, dispersible powders, nondispersible powders, oil solubles, liquid concentrates or aqueous solutions.

The agricultural biocide component can be, for example, a fungicide, herbicide, insecticide, nematocide, acaricide and mixtures thereof. By "agricultural biocide" it is meant any biocide, pesticide, or herbicide. The type of the biocide component used is not critical to the use of the adjuvant composition according to the invention. Examples of common biocides are listed in U.S. Pat. No. 4,793,850 to Koester et al. and in U.S. Pat. No. 4,851,421 to Iwasaki et al., the entire disclosures of which are incorporated herein by reference.

The plant protection composition resulting from the combination of adjuvant and biocide component contains an effective amount of biocide component. The "effective amount" can be experimentally determined and depends upon the type and form of biocide used. For example, a ready-to-dilute plant protecting composition may include from about 1 to about 95% biocide, by weight of the composition. Ready-to-dilute formulations in the dry flowable powder form usually contains greater than about 90% biocide, while other powdered formulations can contain less than 30% biocide. In addition, ready-to-dilute formulations in the form of emulsifiable concentrates can

contain up to about 40% biocide, while the concentration of biocide in suspension concentrates can be about 60%.

The adjuvant composition of the present invention can be added to a tank mix generally in amounts of from .0001 to about 10 % of the tank mix. Preferably, the adjuvant composition is from 0.001% to about 5% by weight of the tank mix. More preferably, the adjuvant composition can be added to the tank mix in amounts of from 0.005% to about 2 % by weight of the formulation. Most preferably, the adjuvant composition is from 0.01% to about 0.5% by weight of the formulation. In a preferred embodiment, the adjuvant composition of the present invention has a Hydrophilic-Lipophilic Balance (HLB) that matches the HLB of the active biocide component. The adjuvant composition of the present invention can be added to the ready-to-dilute biocide formulation generally in amounts of from 1 to 40 percent of the tank mix so as to result in tank mix concentration specified above.

The adjuvant composition of the present invention can also be combined with the biocide component along with other ingredients such as other surfactants and adjuvants. Examples of these optional ingredients include, but are not limited to solvents, co-solvents, preservatives, corrosion inhibitors, thickening agents and buffering agents.

The advantages and other characteristics of the present inventions are best illustrated by the following examples.

EXAMPLES

In the examples, the performance of compositions containing commonly used surfactants were compared with those of compositions containing topped or peaked alcohol ethoxylates. Table I lists the chemical description of the various surfactants used.

TABLE I

	<u>Surfactant</u>	<u>Chemical Description</u>
5	Alfonic 610-50R Alfonic 810-60 Alfonic 1412-60 APSA-80 Genapol UD 079	ethoxylated linear alcohols (50% EO) ethoxylated linear alcohols (60% EO) ethoxylated linear alcohols (60% EO) ethoxylated nonyl phenol C ₁₁ linear alcohol ethoxylate containing an average of 7 moles EO/mole alcohol
10	Genapol 26 L 60 N Genapol UD 070N	peaked C ₁₂₋₁₆ linear primary alcohol containing an average of 7 moles EO peaked C ₁₁ linear alcohol ethoxylate containing an average of 7 moles EO
15	Genapol UD 080N Neodol 1-7	peaked C ₁₁ linear alcohol ethoxylate containing an average of 8 moles EO C ₁₁ linear primary alcohol ethoxylate containing an average of 7 moles EO/mole alcohol
20	Neodol 23-6.5T Neodol 23-9	topped C _{12-C13} linear alcohol ethoxylate containing an average of 6.5 moles EO/mole alcohol C _{12-C13} linear alcohol ethoxylate containing an average of 9 moles EO/mole alcohol
25	Neodol 91-6 Nonoxynol 10 Novel II 1412-70	C _{9-C11} linear primary alcohol ethoxylate containing an average of 6 moles EO/mole alcohol ethoxylated nonyl phenol (10% EO) C _{12-C14} linear primary alcohol ethoxylate (70% EO)
30	Novell II 1012-7 Novel II 1012-5.5	peaked C ₁₀₋₁₂ linear alcohol ethoxylate containing an average of 7 moles EO peaked C ₁₀₋₁₂ linear alcohol ethoxylate containing an average of 5.5 moles EO
35	Silwet Toximul 8304 Triton XL-80N	silicon glycol copolymer alcohol ethoxylate blend containing an average of 6 moles EO/mole alcohol short-chain secondary alcohol alkoxylate blend containing ethylene oxide and propylene oxide
40	X-77	commercial product based on Nonoxynol-10

Following is a list of adjuvants and their manufacturers from which they may be obtained. Alfonic 610-50R, Alfonic 810-60, Alfonic 1412-70, Novel II

1412-70, Novel II 1216CO-7, Novel II 1012-7, Novel II 1012-5.5 can be obtained from CondeaVista. Genapol UD 070, Genapol UD 070, Genapol UD 070N, Genapol 26-L-60N, and Genapol UD 080N can be obtained from Clariant Corporation. Neodol 1-7, Neodol 23-6.5T, Neodol 23-9, and Neodol 91-6 can be obtained from The Shell Chemical Company. Silwet can be obtained from Osi. Toximul 8304 can be obtained from Stepan Chemical Company. Triton XL-80N can be obtained from Union Carbide. Accent Pinnacle can be obtained from DuPont. Beacon can be obtained from Ciba. Roundup can be obtained from Monsanto. Pursuit can be obtained from American Cyanamid. Gramoxone Extra can be obtained from Zeneca.

EXAMPLE 1

Weed control evaluation was performed on sprayable herbicide formulations containing a series of alternate surfactants to replace Nonoxynol-10, the accepted industry standard.

The results of this greenhouse study are shown in Table II. Table II provides the weed control rating in percentage. As shown in Table II, Neodol 23-6.5T, a topped alcohol ethoxylate, out-performed Nonoxynol 10 on the BeaconJ and RoundupJ herbicide brands in Giant Foxtail and Seedling Johnsongrass. Moreover, Neodol 23-6.5T, the topped alcohol ethoxylate was the only alcohol ethoxylate that performed close to Nonoxynol-10.

The amount of herbicide used corresponding to the herbicides is Table II is shown in the immediately following chart.

Chart of Herbicides

Herbicide	Amount
Accent	3.5 g a.i./ha at 25 gpa
Beacon	2.0 g a.i./ha at 25 gpa on seedling Johnson grass and 14.8 g a.i./ha on giant foxtail
Roundup	0.3 lb./acre at 25 gpa
Pursuit	24.7 g a.i./ha at 25 gpa

Pinnacle	1.9 g a.i./ha at 25 gpa
----------	-------------------------

In the above chart, "a.i." means "active ingredient, "ha" is the abbreviation for hectare, and "gpa" means "gallons per acre spray volume."

TABLE II. WEED CONTROL RATING (YEAR 1)

Herbicide	ACCENT	ACCENT	BEACON	BEACON	ROUNDUP	PURSUIT	PINNACLE	
Weed Species	Giant	Seedling	Giant	Seedling	Giant	Velvet leaf	Velvet leaf	
	Foxtail	Johnsongrass	Foxtail	Johnsongrass	Foxtail			Total Average Rating
Adjuvant (at 0.5 %)								
Nonoxynol 10	81	75	66	72	88	71	61	73.4
Neodol 23-6.5T	76	66	71	78	98	43	50	68.9
Neodol 91-6	74	74	53	69	79	39	41	61.3
Neodol 1-7	71	68	65	64	79	36	42	60.7
Genapol UD079	70	72	61	68	76	36	34	59.6
Triton XL-80N	74	73	44	73	79	34	39	59.4
X-77	65	63	58	54	59	53	45	56.7
Toximul 8304	68	55	36	50	83	34	34	51.4
Alfonic 610-50R	61	49	53	56	79	31	28	51.0
Alfonic 810-60	63	43	29	59	73	38	33	48.3
Coupled Ethoxylate	63	34	33	51	64	29	29	43.2
Herbicide Alone	4	13	14	6	61	23	8	18.4
No Treatment	0	0	0	0	0	0	0	0

EXAMPLE 2

Weed control evaluation was performed on sprayable herbicide formulations containing a peaked alcohol ethoxylate (Novel II 1412-70) and compared to formulations containing Nonoxynol-10.

Table III provides the weed control rating in percentage. In this test, herbicides containing the peaked alcohol ethoxylate out-performed those that contained Nonoxynol-10 in weed control experiments on five herbicides and four weed species.

5 The amount of herbicide used corresponding to the herbicides in Table III is shown in the immediately following chart.

Chart of Herbicides

Herbicide	Amount
Accent	3.5 g a.i./ha at 25 gpa
Beacon	2.0 g a.i./ha at 25 gpa on seedling Johnson grass and 14.8 g a.i./ha on giant foxtail
Roundup	0.3 lb./acre at 5 gpa on velvet leaf and lambs quarters, and 0.15 lb. a.i./A on foxtail and Johnson grass
Pursuit	24.7 g a.i./ha at 25 gpa
Pinnacle	1.9 g a.i./ha at 25 gpa

10 In the above chart, "a.i." means "active ingredient, "ha" is the abbreviation for hectare, and "gpa" means "gallons per acre spray volume."

TABLE III. WEED CONTROL RATING (YEAR 2)

Herbicide	Accent	Accent	Beacon	Beacon	Roundup	Roundup	Pinnacle	Pinnacle	Pursuit	Pursuit	Roundup	Roundup	
Weed Specie	Giant Foxtail	Seedling Johnson-grass	Giant Foxtail	Seedling Johnson-grass	Giant Foxtail	Seedling Johnson-grass	Velvet leaf	Common Lambs-quarters	Velvet leaf	Common Lambs-quarters	Velvet leaf	Common Lambs-quarters	Total Average Rating
Adjuvant (at 0.5%)													
Novel II 1412-70	63	70	62	66	95	85	60	78	76	61	88	89	74.4
Nonoxyno I-10	71	71	66	65	88	64	56	76	64	55	68	88	69.3
X-77	53	56	57	44	84	65	44	66	49	50	73	90	60.9
Herbicide Alone	5	4	4	4	88	60	13	13	23	10	73	81	31.4
Untreated	0	0	0	0	0	0	0	0	0	0	0	0	0

5 Table IV contains the weed control rating in percentage. It is surprising that several peaked linear alcohol ethoxylates outperformed non-peaked alcohol ethoxylate, X-77.

TABLE IV. WEED CONTROL RATING (YEAR 3)

	Accent			Pursuit			Pinnacle		Roundup		Gramoxone Extra	Average
	Velvet-leaf	Gt Foxtail	Lambs-quarters	Velvet-leaf	Gt Foxtail	Lambs-quarters	Velvet-leaf	Lambs-quarters	Gt Foxtail	Lambs-quarters	Lambs-quarters	
Novel II 1412-70	50	68	61	58	62	48	80	76	91	71	86	68.4
Genapol 28-L-60N	39	57	53	4	55	44	46	75	78	66	51	55.3
Genapol UD 080N	36	56	43	39	59	43	38	73	65	61	39	50.2
X-77	40	47	51	41	48	43	39	69	46	70	53	49.7
Genapol UD 070N	19	49	46	41	56	46	39	74	58	68	38	48.5
Novel II 1012-7	22	56	46	40	52	50	45	66	58	58	31	47.6
Novel II 1012-5.5	25	47	43	39	48	43	40	74	49	60	31	45.5
Herbicide Alone	10	5	9	25	10	9	17	11	53	52	26	20.6
No Treatment	0	0	0	0	0	0	0	0	0	0	0	0

10

The amount of herbicide used corresponding to the herbicides in Table IV is shown in the immediately following chart. The surfactant comprised 0.25% of the spray volume.

Chart of Herbicides

Herbicide	Amount
Accent	0.03125 lbs/acre at 25 gpa on velvetleaf and lambsquarters and 0.003125 lbs/acre at 25 gpa on giant Foxtail
Gramoxone Extra	0.156 lbs/acre at 25 gpa
Roundup	0.25 lbs/acre at 5 gpa on lambsquarters and 0.1 lbs/acre on foxtail
Pursuit	0.022 lbs/acre at 25 gpa
Pinnacle	0.0016 lbs/acre at 25 gpa

In the above chart, "a.i." means "active ingredient, "ha" is the abbreviation for hectare, and "gpa" means "gallons per acre spray volume."

5 In a total of 74 field trial runs in which a nonoxynol-10 based spray adjuvant was compared with a peaked alcohol ethoxylate formulation, there was no significantly different average weed control between the two formulations. The nonoxynol-10 formulation had a weed control rating of 81.5 % compared to the peaked formulations weed control of 80.5%. Both formulations, the peaked and non peaked, gave weed control that was significantly better than the herbicide alone which had a weed control of 71.8 percent. It is surprising and one would generally expect that crop yield would follow weed control. The surprising result is that the peaked adjuvant formulation had a higher crop yield than the nonoxynol-10 or herbicide treatment alone. The peaked adjuvant formulation had a normalized yield of 106.9%. The nonoxynol-10 formulation gave a normalized yield of 105.3%, which is 1.6% less than the peaked adjuvant formulation. This result is surprising in that the peaked formulation gave comparable weed control to the nonoxynol-10 formulation and gave a higher crop yield over a broad range of crops and herbicides.

Thus, in a preferred embodiment of the present invention, the plant protecting composition comprises a biocide in combination with an adjuvant composition comprised essentially of a nonionic surfactant component selected from the group consisting of topped alcohol ethoxylates, peaked alcohol ethoxylates, topped alcohol propoxylates, peaked alcohol propoxylates, topped propoxylated and ethoxylated alcohols, peaked propoxylated and ethoxylated alcohols and combinations thereof and in combination with conventional alcohol alkoxylates, wherein the peaked surfactants contain less than about 3% residual free alcohol.

Additionally, the plant protecting composition may contain approximately 9% isopropanol, approximately 0.5% sodium benzoate, acetic acid, soft water and other components in addition to the peaked or topped alcohol alkoxylates and combinations thereof in combination with conventional alcohol alkoxylates.

A formulation of one preferred embodiment is shown in the Table V set forth immediately below.

TABLE V. FORMULATION

Compound	Percentage
Toximul 8304 isodecyl Alcohol plus 6 moles of Ethylene Oxide conventional Alcohol Ethoxylate	40
Novell II 1216 C0-7 C _{12, 14} and 16 alcohol with 7 moles EO peaked alcohol ethoxylate	40
Isopropanol	9
Sodium Benzoate	0.5
Acetic Acid	as needed to obtain pH of 7
Soft water	quantity sufficient

Of course, it should be understood that a wide range of changes and modifications can be made to the embodiments described above. It is

therefore intended that the foregoing description illustrates rather than limits this invention, and that it is the following claims, including all equivalents, which define this invention.

WHAT IS CLAIMED IS:

1. A plant protecting composition comprising an agricultural biocide and adjuvant composition comprising a primary component selected from the group consisting of topped alcohol alkoxylates and peaked alcohol alkoxylates and combinations thereof and one or more conventional alcohol alkoxylates.

2. The composition of claim 1 wherein the primary component is a primary aliphatic alcohol alkoxylate.

3. The composition of claim 1 wherein the peaked alcohol alkoxylates are C₆-C₂₀ linear alcohol alkoxylates.

4. The composition of claim 1 wherein the peaked alcohol alkoxylates contain less than about 3% of residual free alcohol.

5. The composition of claim 1 wherein the topped alcohol alkoxylates are linear alcohol alkoxylates having a C₆ or greater alkyl group.

6. The composition of claim 1 wherein the primary component is selected from the group consisting of topped alcohol ethoxylates, peaked alcohol ethoxylates, topped alcohol propoxylates, peaked alcohol propoxylates, topped propoxylated and ethoxylated alcohols, peaked propoxylated and ethoxylated alcohols and combinations thereof.

7. The composition of claim 5 wherein the peaked alcohol ethoxylates comprise from about 2 to about 20 moles of ethylene oxide per mole of alcohol; and

the peaked alcohol propoxylates comprise from about 2 to about 20 moles of propylene oxide per mole of alcohol.

8. A method for increasing the efficacy of an agricultural biocide formulation including a biocide component comprising providing an adjuvant composition comprising a nonionic surfactant component selected from the

group consisting of topped alcohol alkoxylates and peaked alcohol alkoxylates and combinations thereof in combination with a conventional alcohol alkoxylate.

INTERNATIONAL SEARCH REPORT

Inter. Application No
PCT/US 00/00945

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N25/30 A01N57/20 A01N43/40 A01N47/36 A01N43/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 99 26472 A (HENKEL CORP) 3 June 1999 (1999-06-03) the whole document	1-8
X	----- DATABASE WPI Section Ch, Week 199802 Derwent Publications Ltd., London, GB; Class A25, AN 1998-014636 XP002134657 & JP 09 278605 A (AGRO KANESHO KK), 28 October 1997 (1997-10-28) abstract	1-8
X	----- WO 96 00705 A (CHURCH & DWIGHT CO INC) 11 January 1996 (1996-01-11) see: page 9, line 13 - page 10, line 8; Examples I - VI.	1-8
	----- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

3 April 2000

Date of mailing of the international search report

18/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Klaver, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/00945

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 409 885 A (DERIAN PAUL-JOEL ET AL) 25 April 1995 (1995-04-25) column 3, line 54 -column 4, line 13 column 5, line 7-19 ---	1-8
X	WO 91 08666 A (CHEMINOVA AGRO AS) 27 June 1991 (1991-06-27) page 4, line 21-34; examples 1,2 ---	1-8
X	WO 88 09122 A (CHINOIN GYOGYSZER ES VEGYESZET) 1 December 1988 (1988-12-01) page 7, line 3-27; example 13 ---	1-8
X	J. M. GREEN: "Varying surfactant type changes Quizalofop-P herbicidal activity." WEED TECHNOL., vol. 11, 1997, pages 298-302, XP000882667 page 301, column 1, paragraph 1 -column 2, paragraph 4; tables 1-5 ---	1-8
X	J. M. GREEN & J. H. GREEN: "Surfactant structure and concentration strongly affect Rimsulfuron activity." WEED TECHNOL., vol. 7, 1993, pages 633-640, XP000882663 page 638, column 2, paragraph 2 -page 640, column 1, paragraph 3 ---	1-8
X	M. T. GARCIA, I. RIBOSA, J. SANCHEZ LEAL & W. HRECZUCH: "Effect of homolog distribution on the toxicity of alhol ethoxylates." J. AM. OIL CHEM. SOC., vol. 73, no. 7, 1996, XP000882920 see: page 903, col. 1, par. 1 - col. 2, par. 2; 'Results and Discussion'. ---	1-8
X	M. F. COX: "The effect of 'peaking' the ethylene oxide distribution on the performance of alcohol ethoxylates and ether sulfates." J. AM. OIL CHEM. SOC., vol. 67, no. 9, 1990, pages 599-604, XP000882919 see: 'Results and Discussion' ---	1-8
X	K. W. DILLAN: "Effects of the ethylene distribution on nonionic surfactant properties." J. AM. OIL CHEM. SOC., vol. 62, no. 7, 1985, pages 1144-1151, XP000882921 the whole document -----	1-8

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 00/00945

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9926472 A	03-06-1999	AU 1373999 A	15-06-1999
JP 9278605 A	28-10-1997	NONE	
WO 9600705 A	11-01-1996	US 5389386 A AU 2603795 A US 5432147 A	14-02-1995 25-01-1996 11-07-1995
US 5409885 A	25-04-1995	FR 2678807 A CA 2073480 A EP 0522891 A FI 923143 A JP 5186301 A NO 922681 A	15-01-1993 10-01-1993 13-01-1993 10-01-1993 27-07-1993 11-01-1993
WO 9108666 A	27-06-1991	DK 649089 A AT 102791 T AU 649646 B AU 7045791 A CA 2068646 A CS 9006339 A DE 69007494 D EP 0506788 A FI 922808 A HU 62180 A JP 5504562 T LV 10146 A NO 173721 C NZ 236579 A PH 27393 A PT 96261 A TR 24984 A US 5118338 A ZA 9010192 A	21-06-1991 15-04-1994 02-06-1994 18-07-1991 21-06-1991 15-09-1991 21-04-1994 07-10-1992 17-06-1992 28-04-1993 15-07-1993 20-10-1994 26-01-1994 26-03-1993 21-06-1993 30-09-1991 01-09-1992 02-06-1992 30-10-1991
WO 8809122 A	01-12-1988	HU 46836 A DE 3872356 A EP 0317582 A JP 1503304 T RU 2020819 C TR 27149 A US 5045311 A	28-12-1988 30-07-1992 31-05-1989 09-11-1989 15-10-1994 09-11-1994 03-09-1991